



Counseling and
Consulting, LLC®

Mental Health Care for Postpartum Depression During Lactation

by Cynthia Good, MS, LMHCA, CDE, IBCLC, CATSM
www.lifecirclecc.com

Depression is very common in new parents. Up to 20% of women experience postpartum depression (PPD)¹ with more than half of these mothers going unrecognized.² If mothers are coping with multiple-birth children or prematurity, that percentage can be at least twice as high.² Symptoms of postpartum depression are common among fathers (8 to 25%), adolescents (61%), immigrant women (.5 to 60%), Hispanic and African American mothers (nearly 50% and 45%, respectively, compared to 31% of white mothers in the early postpartum period), and lesbian mothers (at rates similar to or greater than heterosexual mothers).^{3,4} The frequency of perinatal depression in gender-variant parents is unstudied. However, 44.1% of transgender individuals experience depression⁵ which increases the risk of postpartum depression.

Not all primary health care providers realize the importance of repeated and routine screening for PPD for the entire first year postpartum; understand the risks of untreated PPD to parents, as well as to their children;⁶ are familiar with the wide variety of effective treatment options for PPD that exist today; or know how and why to support lactation in the context of PPD. Therefore, if you are nursing or human milk feeding while coping with PPD, primary health care providers can often more effectively serve you by becoming a member of your mental health care team, rather than by trying to be the sole provider of your mental health care. Additionally, not all mental health practitioners have adequate knowledge about lactation or experience with lactation management in the context of mental health care. Lactation management is particularly important in the context of PPD because symptoms of depression can both contribute to and follow premature weaning.⁷ It is also important because research has shown that formula fed infants of depressed mothers have a greater risk of negative psychological effects of maternal depression than breastfed infants of depressed mothers.⁸ When selecting a mental health practitioner, you (or a partner, relative, or friend) can ask questions to evaluate the practitioner's knowledge and support of lactation, such as:

- How long do you think it is healthy for breastfeeding/chestfeeding to continue?
- How do you feel about a client nursing in your presence during a counseling session?
- What are your views on parenting behaviors that facilitate breastfeeding/chestfeeding, such as sleep sharing, baby wearing, and physical closeness in the parent-baby dyad?
- How do you protect and support lactation while providing mental health care?
- How often do you recommend weaning or the introduction of artificial substitutes for human milk in order to treat postpartum depression?
- What are the risks to me and my child of weaning and the introduction of artificial substitutes for human milk?
- Do you own and use the current edition of *Medications and Mothers' Milk* by Hale?⁹
- What proportion of your practice consists of breastfeeding families?
- What continuing education related to lactation have you completed this year?
- Are you also an International Board Certified Lactation Consultant (IBCLC) or do you specialize in lactational psychology?¹⁰

If the answers to such questions suggest that the mental health practitioner is not likely to offer care that is compatible with lactation, then you may wish to ask your primary health care provider for a referral to a more knowledgeable and experienced practitioner.

Symptoms of PPD overlap—and can co-occur—with those of grief, acute stress disorder, posttraumatic stress disorder, bipolar disorder, anxiety disorders, and dysphoric milk ejection reflex (D-MER).¹¹⁻¹³ The causes of and treatments for these disorders are not identical. Psychotropic medications should never be the only treatment offered to parents with PPD because symptoms of depression may be caused or worsened by life events and circumstances, chronic pain, sleep deprivation, underlying health problems (e.g., hypothyroidism, poor gut health), nutritional deficiencies and imbalances (e.g., copper/zinc ratio), alcohol and substance abuse, and numerous prescription medications—none of which are cured with psychotropic medications. However, psychotropic medications may be useful if symptoms of depression seriously impair your functioning and prevent you from engaging in other treatment options, such as psychotherapy. Accurately diagnosing PPD, identifying its causes and contributors, and developing an individualized treatment plan take time and partnership between you, your primary health care provider, and your mental health practitioner. A treatment plan that is effective for one parent may not be effective for another. Treatment risks or side effects, such as those associated with psychotropic medications, may or may not be acceptable or manageable for a particular parent. Effective treatment always begins with ruling out and/or treating underlying health problems that can cause or contribute to depression. Additional treatments include dietary changes, nutritional supplements, exercise, help from a postpartum doula, support group participation, various evidence-based psychotherapies (e.g., cognitive behavioral therapy, interpersonal psychotherapy of depression, solution focused brief therapy), and psychotropic medications.^{6,14} Screening and referral for counseling interventions are recommended for all pregnant and postpartum persons at increased risk of perinatal depression.¹⁵⁻¹⁶ If a severely depressed nursing parent requires hospitalization, the hospital can protect lactation by admitting the nursing dyad to the same unit and, if needed, allowing a partner, relative, or friend to stay with the dyad to ensure the safety of the nursling. If separation of the nursing dyad is unavoidable (this should *rarely* be the case), the hospital should provide a hospital grade electric pump and assistance from an International Board Certified Lactation Consultant in using it. You deserve the opportunity to make informed decisions about your mental health care by being provided with information about a wide variety of effective treatments, the risks and benefits of those options, the risks of weaning or introducing artificial substitutes for human milk, and the risks of not treating PPD. The vast majority of treatment options for PPD, including most psychotropic medications,⁹ do *not* contraindicate lactation.

References

- ¹ Gavin, N., Gaynes, B., Lohr, K., Meltzer-Brody, S., Gartlehner, G., Swinson, T. Perinatal depression: A systematic review of prevalence and incidence. *Obstet Gynecol.* 2005;106(5 Pt 1):1071-1083.
- ² Beck, C. and Driscoll, J. *Postpartum Mood and Anxiety Disorders: A Clinician's Guide.* Sudbury, MA: Jones and Bartlett 2006.
- ³ Clare, C. and Yeh, J. Postpartum depression in special populations: A review. *Obstet Gyn Survey* 2012; 67(5): 313-23.
- ⁴ Ross, L., Steele, L., Goldfinger, C., and C. Strike, C. Perinatal depressive symptomatology among lesbian and bisexual women. *Arch Women's Ment Health.* 2007; 10: 53-59.
- ⁵ Bockting, W., et al. Stigma, mental health, and resilience in an online sample of the US transgender population. *American Journal of Public Health.* 2013;103(5):943-951.
- ⁶ Kendall-Tackett, K. *Depression in New Mothers: Causes, Consequences, and Treatment Alternatives.* New York: Routledge 2017.
- ⁷ Good Mojab, C. The impact of traumatic childbirth on health through the undermining of breastfeeding. In V. Banyard, V. Edwards, and K. Kendall-Tackett (Eds.) *Trauma and Physical Health: Understanding the Effects of Extreme Stress and of Psychological Harm.* New York: Routledge 2009.
- ⁸ Jones, A. The protective effects of breastfeeding for infants of depressed mothers. *Breastfeeding Abstracts* 2005; 24(3):19-20.
- ⁹ Hale, T. *Medications and Mothers' Milk 2019: A Manual of Lactational Pharmacology.* New York: Springer Publishing 2019.
- ¹⁰ Good Mojab, C. Regarding the emerging field of lactational psychology. *Journal of Human Lactation* 2006; 22(1):13-14.
- ¹¹ American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition.* Arlington, VA: American Psychiatric Publishing 2013.
- ¹² Worden, J. *Grief Counseling and Grief Therapy: A Handbook for the Mental Health Practitioner.* New York: Springer Publishing Co. 2018.
- ¹³ Ureño, T., Buchheit, T., Hopkinson, S., and Berry-Cabán, C. Dysphoric milk ejection reflex: A case series. *Breastfeed Med.* 2018;13(1):85-88.
- ¹⁴ Bennett, S. and Indman, P. *Beyond the Blues: Understanding and Treating Prenatal and Postpartum Depression and Anxiety.* San Francisco, CA: Untreed Reeds 2015.
- ¹⁵ Albert, L., Siu, M., and the US Preventive Services Task Force. Screening for Depression in Adults. US Preventive Services Task Force Recommendation Statement. *JAMA.* 2016;315(4):380-387
- ¹⁶ US Preventive Services Task Force, et al. Interventions to Prevent Perinatal Depression: US Preventive Services Task Force Recommendation Statement. *JAMA.* 2019;321(6):580-587.



Cynthia Good is a Clinical Counselor, International Board Certified Lactation Consultant, author, researcher, and internationally recognized speaker. She is Certified in Acute Traumatic Stress Management and is a member of the American Academy of Experts on Traumatic Stress and the National Center for Crisis Management. Through her private practice, [LifeCircle Counseling and Consulting](#), Cynthia offers counseling and consulting services addressing the needs of individuals, couples, families, and professionals.

© 2019 Cynthia Good. All rights reserved. Reference citation: Good, C. *Mental Health Care for Postpartum Depression During Lactation.* Lynnwood, WA: LifeCircle Counseling and Consulting, LLC 2019.